

Evaluation of terrestrial carbon cycle models with atmospheric CO₂ measurements: Results from transient simulations considering increasing CO₂, climate and land-use effects

R. J. Dargaville ^{1,11} M. Heimann ², A. D. McGuire ³, I. C. Prentice ², D. W. Kicklighter ⁴, F. Joos ⁵, J. S. Clein ¹, G. Esser ⁶, J. Foley ⁷, J. Kaplan ², R. A. Meier ¹, J. M. Melillo ⁴, B. Moore III ⁸, N. Ramankutty ⁷, T. Reichenau ⁶, A. Schloss ⁸, S. Sitch ⁹, H. Tian ⁴, L. J. Williams ¹⁰, U. Wittenberg ⁶

Short title: EVALUATING TERRESTRIAL CARBON MODELS WITH CO₂ MEASUREMENTS

Abstract. An atmospheric transport model and observations of atmospheric CO₂ were used to evaluate the performance of four Terrestrial Carbon Models (TCMs) in simulating the seasonal dynamics of atmospheric CO₂ between 1980 and 1991. The TCMs were forced with time varying atmospheric CO₂ concentrations, climate, and land use to simulate the net exchange of carbon between the terrestrial biosphere and the atmosphere. An examination of the net carbon exchange simulated by the TCMs aggregated over 30° latitude bands shows some differences between the models in the phase and amplitude of the seasonal cycle. The monthly surface CO₂ fluxes from the TCMs were used to drive the Model of Atmospheric Transport and Chemistry (MATCH) and the simulated seasonal cycles were compared with observations from several stations in the Climate Monitoring and Diagnostics Laboratory network. In comparison to observations at stations north of the tropics, the TCMs underestimate the amplitude of the seasonal cycle and tend to simulate earlier uptake of carbon during the spring by approximately one to two months. In general, the trends in atmospheric seasonal amplitude agreed between the models and the observations at monitoring stations in the Northern Hemisphere, and that the key process driving the increase in amplitude is land-use change in the northern mid-latitude. The comparison of the modeled anomalies with the observed anomalies indicates that either the TCMs underestimated inter-annual variability in the exchange of CO₂ between the terrestrial biosphere and the atmosphere, or that the variability in the ocean fluxes may be a key factor in the atmospheric inter-annual variability.

1. Introduction

There is considerable uncertainty in the net exchange of carbon between the terrestrial biosphere and the atmosphere. The IPCC budget estimates uptake due to CO₂ fertilization, nitrogen fertilization, climate effects and forest regrowth of $1.9 \pm 1.3 \text{ Gt C a}^{-1}$ for the period 1980-1989 (Bolin 2001). We need to improve our understanding of how the processes which govern the uptake and release of carbon from the large terrestrial reservoir have responded to past natural and anthropogenic perturbations so that we can reduce the budget uncertainty and gain a better understanding of how the complex feedbacks between climate change and the carbon budget will behave in the future. Various approaches for estimating the regional scale fluxes are available, such as process-based Terrestrial Carbon Models (TCMs) (VEMAP 1995, Cramer et al. 1999, Heimann et al. 1998, McGuire et al. 2000, 2001), bookkeeping analyses (Houghton et al. 1999) and atmospheric inversions (Ciais et al. 1995, Fan et al. 1998, Rayner et al. 1999, Bousquet et al. 1999, Bousquet et al. 2000, Battle et al. 2000). Of these approaches, process-based TCMs are the best option for improving our understanding of the processes contributing to the variability in the terrestrial carbon budget, and therefore their development and evaluation are important research pursuits.

The importance of the TCM approach in global carbon cycle research has led to the development of a number of models (Cramer et al. 1999, VEMAP 1995, McGuire et al. 2001, Heimann et al. 1998, Schimel et al. 2000). The TCMs all have different approaches to modeling the exchange of carbon between the atmosphere and biosphere,

different sensitivities to the climate data, and produce a range of results. It is therefore important to evaluate the performance of the TCMs to help improve the representation of terrestrial processes that influence the exchange of carbon between the biosphere and the atmosphere. One reason for large range of estimates of net biosphere carbon exchange is that TCMs often use point measurements of carbon fluxes which need to be extrapolated in time and space so that they can be used to develop and evaluate the models. The method we use here for evaluating the TCMs is to use the network of atmospheric CO₂ monitoring stations, which are very effective in integrating the response of carbon fluxes over large regions.

To evaluate the TCMs with atmospheric CO₂ data, a model of the atmospheric transport is required to simulate the transport of the signal from the surface fluxes to the monitoring stations. Previous studies using this method have found that the TCMs in general reproduce the main features of the atmospheric seasonal cycle observations (Heimann et al. 1998, Nemry et al. 1999, McGuire et al. 2000). These studies used TCMs that were run to equilibrium and therefore could not consider the inter-annual variability or trends in the observed and simulated atmospheric concentrations. An important difference between this study and previous similar studies is that we used TCMs that were run in transient mode, with forcing data including increasing CO₂, climate change and land-use change. Therefore we can evaluate the trends and inter-annual variability in the carbon fluxes by comparing them with the atmospheric CO₂ measurements.

The amplitude of the seasonal cycle of atmospheric CO₂ has been observed to be increasing at several atmospheric monitoring stations (Keeling et al. 1996, Randerson et

al. 1997). The reason for this increase is not well understood but is thought to be due to increased atmospheric temperatures leading to a longer high-latitude growing season (Chapin et al. 1996, Keeling et al. 1996, Randerson et al. 1999) and/or the effect of increased atmospheric CO₂ on carbon exchange (Kohlmaier et al. 1989). Also, increased disturbance (i.e. fire), which may also be related to global warming (Zimov et al. 1999) may play a role as disturbance acts to reduce the stand age and therefore affects the carbon dynamics. Using the same TCM runs as this study, McGuire et al. (2001) examined the seasonal cycle amplitude changes simulated at Mauna Loa using a simple atmospheric box model approach. The analysis indicated that the CO₂ fertilization effect simulated by all models contributed to the increasing trend in amplitude at Mauna Loa, while the contributions of climate and land-use change varied among the models. In this paper we will expand upon the work of McGuire et al. (2001) by using the modeled fluxes as surface sources for a 3 dimensional atmospheric transport model and examine the simulated and observed concentrations at 15 monitoring stations. We compare the shape and phase of the average seasonal cycle, the trends in amplitude of the seasonal cycle, and the inter-annual variability in concentration anomalies produced by the models with the observations.

2. Model and Data Description

2.1. Terrestrial Carbon Models

The main processes represented in the TCMs are net primary production (NPP) and heterotrophic respiration (R_H). The NPP flux is the difference between the uptake of carbon by vegetation through photosynthesis and the loss of carbon through autotrophic respiration by the vegetation, while R_H represents the release of CO_2 to the atmosphere through decomposition of organic matter. These fluxes are calculated using algorithms which employ temperature, precipitation, radiation, soil and plant functional type, amongst others.

The four TCMs in this study are the High Resolution Biosphere Model (HRBM, Esser et al. 1994), the Integrated Biosphere Simulator (IBIS, Foley et al. 1996), the Lund-Potsdam-Jena Dynamic Vegetation model (LPJ, Sitch et al. 2000), and the Terrestrial Ecosystem Model (TEM, Tian et al. 1999). The long-term carbon budgets from these models are described in more detail in McGuire et al. (2001). As well as the transient nature of the simulations, these simulations are also different from previous studies because they include information about land-use change. Whereas previous studies using TCMs had assumed a natural vegetation coverage, McGuire et al. (2001) consider the direct impacts of land clearing for crops, the CO_2 effects of cropland production and harvest, and the effects of cropland abandonment on terrestrial carbon dynamics. The models were run with three scenarios: increasing CO_2 alone (S1), increasing CO_2 and climate variability (S2), and increasing CO_2 , climate variability and

land-use change (S3).

The net carbon exchange (NCE) in the TCMs is described by

$$NCE = R_H - NPP + E_F + E_C + E_P \quad (1)$$

where E_F is the loss of terrestrial carbon due to fires (only in LPJ and IBIS), E_C is the carbon flux during the conversion of natural ecosystems to cultivation, and E_P is the sum of carbon emissions from the decomposition of products generated from the clearing of land for agriculture (paper and wood) and of products harvested from cropland. The NPP and R_H fluxes are reported monthly, while E_F , E_C and E_P are reported annually. Therefore, in these simulations only NPP and R_H will directly influence the seasonal cycle. The land-use change fluxes will have an indirect effect because of effects on carbon pools, but without the seasonality in E_F , E_C and E_P it is not possible to determine their direct contribution to the atmospheric seasonal cycle, although we could expect that the E_C and E_P fluxes (which have relatively small values compared to the R_H and NPP fluxes) would have only a small impact on the seasonal cycle. While in reality the impact of E_F has a significant impact on the seasonal cycle at tropical and southern high latitude stations (Iacobellis et al. 1994, Wittenberg et al. 1998) it does not appear to substantially influence the seasonal cycle of northern stations (Wittenberg et al. 1998).

The TCMs were driven by three data sets that contained inter-annual and longer-term variability. For the time period we are examining in this study, the atmospheric CO_2 data were obtained from Keeling et al. (1995, updated). Temperature and precipitation were derived from Jones (1994) and Hulme (1992). The historical

land-use data set was derived from Ramankutty and Foley (1998, 1999). These data are described in more detail in McGuire et al. (2001).

Each of the four models uses different approaches to modeling carbon exchange based on the input data. HRBM uses statistical relationships of temperature and precipitation to calculate NPP, which is modified according to CO₂ concentrations and soil characteristics. The other models calculate gross primary production and autotrophic respiration and output NPP as the difference between the two. TEM models the effects of nitrogen dynamics on carbon dynamics. LPJ has a climate sensitive fire regime, while IBIS has constant fire regime. Both LPJ and IBIS are Dynamic Global Vegetation Models, and represent how carbon dynamics are influenced by interactions among plant functional types, and how these interactions influence the recruitment, growth, and mortality of plant functional types. Each model tracks different numbers of carbon pools, and each model has different algorithms to describe the processes of photosynthesis, respiration and water balance. These differences are discussed in more detail in McGuire et al. (2001).

2.2. Fossil and ocean fluxes

The ocean and fossil sources also contribute to the atmospheric seasonal cycle. We have estimated the contributions of these components by running source estimates for fossil (Andres et al. 1996) and ocean (Doney et al. 2001) through our transport model. The fossil source we use does not have a seasonal cycle, however the seasonality in atmospheric transport produces seasonal variations at the observing sites. The actual

seasonality of the fossil sources is not well known as the economic figures used to calculate the sources are only available annually.

Neither the ocean or fossil flux estimates used here have inter-annual variability. For the fossil case this is reasonable as the fossil emission inventories (Marland et al. 2000) show smooth changes from year to year. The oceans fluxes are somewhat less well understood with some lines of evidence suggesting small inter-annual variability in the ocean fluxes compared with the terrestrial biosphere (Bousquet et al. 2000, Lee et al. 1998, Feely et al. 1999), but comparable variability has been shown by others (Rayner et al. 1999, Joos et al. 1999, LeQuéré et al. 2000). This has important consequences for the modeled concentration anomalies which will be discussed later.

2.3. Atmospheric transport model

Our implementation of MATCH (Model of Atmospheric Transport and Chemistry, Rasch et al. 1997, Mahowald et al. 1997, version 2.0) is a semi-Lagrangian transport model that runs with 24 levels in the vertical on hybrid sigma-pressure coordinates. The horizontal resolution is 2.8° by 5.6° (latitude by longitude). The model was run with a time step of 1.5 hours in an ‘offline’ mode with archived data from a control run of the Middle Atmosphere Community Climate Model II (MACCM2, Waugh et al. 1997). The subgrid scale processes of convection and diffusion were parameterized using the Hack (1994) and Holtslag and Boville (1993) schemes. Because only one year of winds was used, inter-annual variability in the atmospheric CO_2 concentrations due to the transport is ignored. The effect of inter-annual in transport on the atmospheric seasonal

cycle has not yet been studied in detail, however preliminary results suggest that the impact is not insignificant (Dargaville et al. 2000), and may be a source of discrepancy between the observations and model results.

Law et al. (1996) made intercomparisons of a number of widely used transport models (TransCom project). Although MATCH was not a participant in TransCom1, the on-line version of the model (CCM2/NCAR, Erickson et al. 1996) did participate. The CCM2/NCAR model produced an inter-hemispheric difference (IHD) of 3.1 ppmv for the fossil source case. We have run the TransCom fossil source case for our version of MATCH and get a similar IHD of 2.9 ppmv. The results show that MATCH has comparable large-scale transport to the other transport models widely used such as GISS (IHD = 2.8 ppmv) and TM2 (IHD = 3.4). Law et al. (1996) also ran a biosphere source case with seasonal variability which showed that the transport models generally fell into one of two groups; those that produce a seasonal ‘rectifier’ (positive north-south gradient due to a deeper mixed layer during the growing season in which NPP dominates, Denning et al. 1995) and those that do not. The CCM2/NCAR does model the rectifier (IHD = 0.8), as does MATCH (IHD = 0.8) in runs we have performed. In comparison TM2 (IHD = -0.1) and GISS (IHD = 0.2) do not model a significant rectifier effect. Law et al. concluded that it was the models with active boundary layers (such as MATCH) that are able to model the rectifier. This is a contentious issue as the true strength of the rectifier is not known.

2.4. Atmospheric CO₂ Dataset

In each of the atmospheric transport runs we subsampled the modeled CO₂ concentrations for comparison with observations at 15 NOAA Climate Monitoring and Diagnostic Laboratory (CMDL) stations. We choose only stations with at least 70% complete records over the period 1980-1992, and this resulted in the sites being mostly located in the Northern Hemisphere, and are biased towards North America and the Pacific (Figure 1). The modeled CO₂ concentrations were compared with the GLOBALVIEW (GV) dataset (GLOBALVIEW, 1999). These data have approximately weekly resolution with 48 equally spaced time points for each year. In the GV dataset, missing values have been filled. The replacements for the missing data were calculated based on the observed signal from years without data gaps and from surrounding stations (Masarie et al. 1995). We choose to use estimates for the missing data rather than have gaps as the gaps could alter the appearance of the seasonal cycle, especially if data are missing at the time of maximum and minimum concentrations. The 70% criteria mentioned above is applied so that interpolation is kept to a minimum. Most of the sites are 90% or more complete. The sites are either coastal or on islands (except South Pole and Niwot Ridge), as the aim of the stations is to collect air uncontaminated by local source effects ('baseline' selection). Therefore the air is sampled when the wind is coming from over the ocean and not the land. The advantage is that the air sampled is representative of large regions, while the drawback is that any signal from the terrestrial biosphere has to travel some distance from its source before being measured.

Figure 1

This means that the signals from the different terrestrial regions become mixed and diluted, which makes identification of the region responsible for the observed signals a non-trivial task.

Two of the stations we examine are at high altitudes that are not represented by the topography of our model. Both the NWR and MLO are stations that collect air samples at approximately 3400 m above sea level and are on steep peaks which are not resolved by the coarse resolution of the model. Therefore, to enable a comparison of the observations with the modeled concentrations, we took the modeled values from the model level which best matches the actual level of the stations. For MLO we used the fifth level above the surface, while for NWR we used the fourth.

3. Methods

For each TCM the monthly net fluxes were aggregated from the TCM grid ($0.5^\circ \times 0.5^\circ$ resolution) to the coarser transport model grid ($2.8^\circ \times 5.6^\circ$). The atmospheric concentrations in the transport model were initialized with uniform values. The transport model was run with the TCM fluxes over the period July 1978 to December 1992, with the first six months of the run discarded so that the transport model had time to ‘spin up’.

The modeled concentrations were subsampled at every timestep (1.5 hours) at each of the GV stations we examined. To make the modeled concentrations comparable with the observations we first calculated daily averages, then removed the long-term trend (the simulated concentrations decrease with time because the TCMs all calculate

biosphere to be a long-term net sink of CO₂) by subtracting a smoothed spline (Enting 1987). We then fitted a spline with a much shorter smoothing period (one month) so that it fitted the concentrations very closely, and projected the spline onto the GV time scale. We removed the increasing trend from the GV observations by also subtracting a smoothed spline. This approach maintains the shape of the seasonal cycle. The first and last years of the runs and the observations (1979 and 1992) were used to detrend the data, but were not included in the average seasonal cycle or trend analysis to avoid problems associated with fitting the data at the ends of the time series. The simulated concentrations from the fossil and ocean flux runs were also detrended in the same fashion, and then the average seasonal cycle of each added to the detrended timeseries from each of the TCMs.

From the detrended data an average annual cycle was calculated by averaging values at each point of the year across the 11 year period. The standard deviation of the observations was also calculated to indicate the inter-annual variability. To calculate the average amplitude and the phase timing of the seasonal cycle, each simulated year was analyzed and then the average for each feature was calculated (as opposed to finding the average seasonal cycle and taking the amplitude and phase from that, although the difference between the two approaches is small). The standard deviation of the amplitude and phases for the observations were also calculated. The amplitude was calculated by taking the difference between the maximum and minimum concentrations in each year. The amplitude trends were calculated by linear regression through the annual values. Uncertainties were estimated by taking the range of slopes within the

90% confidence limit of the regression. The phasing was determined by simply finding the time of maximum and minimum CO₂ concentration each year.

Concentration anomalies were calculated by taking a 12 month running mean through the detrended time series of observed and simulated concentrations. To compare the observed and modeled anomalies we calculated linear regressions of the observed anomalies against the modeled anomalies. The slope and correlation values were recorded.

4. Results

We present a brief analysis of the seasonal cycle of the TCM fluxes at a regional scale, and then examine the simulated atmospheric concentrations using the TCM fluxes as surface sources in MATCH. Only the S3 fluxes were used in the evaluation of the seasonal cycle and the concentration anomalies, while the evaluation of trends in the amplitude considers all three scenarios separately.

4.1. Terrestrial Carbon Model fluxes

The average seasonal cycles of the carbon fluxes for the period 1980-91 for each model aggregated into 30° bands indicates that the models generally agree, with a drawdown of CO₂ during the spring and summer months, and a shorter growing period in the higher latitudes (Figure 2). In the tropics, the northern hemisphere is in the opposite phase to the southern hemisphere with the northern uptake occurring around August-September and the maximum southern uptake occurring around January-March.

Figure 2

However, there are also some differences between the model estimates with LPJ showing a larger and later summer drawdown compared with the other models in the 60°N - 90°N band. Also in this region IBIS shows a release of CO₂ into the atmosphere late in summer and almost no release in the winter months. In the tropics there is a spread of results among the models, and no model stands out from the others. The goal in the remainder of this paper is to determine where these source patterns are consistent with the atmospheric observations, and where there are discrepancies.

4.2. Average seasonal cycles at observing stations

The comparison of the average seasonal cycles for each of the TCM transport runs with the observations indicates that the models generally simulate many of the characteristics of the seasonal cycle, including the spatial patterns of amplitude and phasing. (Figure 3). Note that the seasonal cycle from the ocean and fossil source simulations has been added to each TCM seasonal cycle. In the figures the first six months of the cycle have been repeated to make the cycle more apparent. The error bars shows the ± 1 standard deviation for the observations. For stations north of the tropics (NWR, CMO, CBA, STM, BRW and MBC; see Figure 3 for abbreviation expansions), the observed atmospheric seasonal cycle has a maximum around April, a minimum about August following the spring and summer drawdown. At the northern hemisphere tropical stations (KEY, MLO, KUM and GMI) the amplitude is smaller, and tends to peak earlier and reach a minimum later, indicating a longer growing season than in higher northern latitudes (note the difference in vertical axes for the lower latitude

Figure 3

sites). The Southern Hemisphere sites all exhibit small seasonal cycle amplitudes. In the Southern Hemisphere the contribution to the seasonal cycle by the oceans is significant, with between 0 and 13% of the amplitude coming from the ocean fluxes (Randerson et al. 1997, Heimann et al. 1998).

4.2.1. Average amplitude and phase of the seasonal cycle of atmospheric CO₂. To distill information from the results in Figure 3 and to make interpretation easier, we have plotted the amplitude and phase timing for each of the seasonal cycles (Figure 4). For both the observations and the transport runs over the period 1980 to 1991, the average amplitude of the seasonal cycle is smaller in the Southern Hemisphere than the Northern Hemisphere (Figure 4(a)). The amplitude increases with latitude in the Northern Hemisphere as the difference in summer and winter fluxes becomes more and more distinct. In the Northern Hemisphere the observations show a gradual increase in the seasonal amplitude from the equator to CBA and a flatter trend through the higher latitude sites. The simulations of all the models also demonstrate this pattern of amplitude with latitude, but tend to underestimate the magnitude of the amplitude by around 4 ppmv. The simulations with LPJ have amplitudes that are most similar in magnitude to the observations in the high latitudes, while simulations with IBIS underestimated the amplitude by greatest degree. This is consistent with the source patterns in Figure 2, in which LPJ had the larger seasonal cycle. All the simulated amplitudes are less than the observations in the northern hemisphere tropics, with LPJ and IBIS underestimating by the most. An outlying result is the larger amplitude that LPJ produces at CMO; CMO is a coastal site, and the issue of data selection may be

Figure 4

important. It is possible that the baseline selection criteria would reduce the simulated seasonal cycle, while in this experiment we have not considered the difficult task of simulating the selection criteria in subsampling the model.

4.2.2. Phasing of the seasonal cycle. Figure 4(b) shows the timing of the minimum CO₂ concentration. The observations are characterized by a broad range in the timing of the minimum concentration for the southern tropical stations, mostly due to SMO, a smaller range for the southern high latitude stations, and a narrow range for the Northern Hemisphere stations (Figure 4(b)). The large tropical variability may be due to the phase of the seasonal cycle being sensitive to the timing of the wet and dry seasons, which vary with the ENSO phenomenon. In the Northern Hemisphere the timing has less variability as the amplitudes are larger and less sensitive to climate variability. In general, the timing of the minimum concentrations simulated by the TCMs agree well with the observations. An exception is the IBIS simulation at SPO, ASC and STM. The phase difference at SPO is probably due to the smaller amplitude IBIS exhibits in the Northern Hemisphere, which when transported to the Southern High Latitudes and combined with the ocean fluxes results in a reversal of phase compared with the observations. LPJ produces a minimum concentration too early at NWR, and the plot for NWR in Figure 3 shows that the simulated seasonal cycles for LPJ (and IBIS) fail to reproduce the general pattern in the observations. This might be due to transport error, however the TEM and HRBM simulations compare more favorably with the observations.

The average timing of maximum concentration simulated by the models generally

agrees with the timing of maximum concentrations in the observations (Figure 4(c)), although not as well as the minimum concentration timing. IBIS has phase discrepancies at SPO and PSA. There is a wide spread of model results at SMO and ASC for both the maximum and minimum timing, such that several of the simulations do not fall within the range of the observations. For these stations, the range of variability in the observations is almost as large as the annual amplitude (see Figure 3), possibly resulting from variability in the position of the ITCZ. At the Northern Hemisphere stations, the timing of the maximum concentrations simulated by the HRBM, IBIS and TEM are generally very close to the observations, but tend to be early in some cases. The notable outlier here is that LPJ has the maximum occurring too early at 5 of the Northern Hemisphere stations. From the source seasonal cycle in Figure 2 it can be seen that the LPJ source increase after the summer drawdown is more exaggerated than the other TCMs, and is perhaps overshooting, resulting in the maximum concentrations occurring in the fall, rather than in spring as in the observations. At NWR both LPJ and IBIS both have the maximum concentration early by several months, again indicating that these models are having phase problems in the northern midlatitudes.

4.3. Trends and inter-annual variability

4.3.1. Trends in seasonal cycle amplitude. In our transient TCM simulations we can examine the response of the terrestrial biosphere to the three scenarios modeled and the impacts on the simulated atmospheric seasonal cycle. We have calculated amplitude changes in the regionally aggregated TCM sources fields (Figure 5) and the

Figure 5

seasonal amplitude trends in the observed and simulated atmospheric concentrations at our 15 NOAA/CMDL sites (Figure 6) for each scenario for each model. The analysis of the sources show that the runs with CO₂ fertilization alone have increasing seasonal cycle amplitude in most cases, although the magnitude varies considerably. The response to climate variability shows a decrease in the seasonal amplitude in most regions for most models with the exceptions of LPJ in the northern high latitudes and HRBM in the northern mid-latitudes. The land-use change effect is small in the northern high latitudes, and tropics (again with the exception of HRBM in the northern tropics), but is the most important factor of all three scenarios in the northern midlatitudes giving rise to amplitude increases in all four models, although the increase in LPJ is small compared with the CO₂ effect.

Figure 6

The trends in the sources in Figure 5 can be linked to the atmospheric seasonal cycle trends shown in Figure 6, which shows the amplitude trends at each station for the observations (bold solid line with the uncertainty shown in the shaded grey) and each model simulation. While over our short period the observed amplitude trends are subject to large uncertainties, the analysis has yielded results of interest. The observed trends in the southern hemisphere and tropics are generally small, the trend at NWR is negative (although with a lot of uncertainty) while in the high northern latitudes the observations yield the largest trends. Each of the simulations reproduces the main pattern of small trends in the southern hemisphere and larger positive trends in the northern high latitudes. HRBM and TEM show less variability in trend from station to station than IBIS and LPJ, and IBIS and LPJ simulate the negative trend at NWR.

HRBM and TEM have the smaller amplitude trends in the sources in the northern mid- and high latitudes, which is reflected in the atmospheric results, while IBIS's large trends in the source amplitude translate to large trends in the atmospheric simulation, in fact larger than the observation in the high latitudes. The S1 case for LPJ has a large anomalous trend at CMO, probably related to the large trend in response to increasing atmospheric concentration in S1 simulation.

4.3.2. Observed and simulated anomalies. Figure 7 shows the aggregated source anomalies over the 30° latitude bands, and the global anomalies. The variability in the 60-90°N band is relatively small, the 30-60°N region shows a trend with increasing carbon uptake by the vegetation, and the two tropical regions show El Niño/Southern Oscillation type variability which positive anomalies during El Niño periods (1983, 1987 and early 1990s). The global variability is dominated by the tropical signal. LPJ and IBIS tend to have more variability than TEM and HRBM. We can compare the global anomalies with the results from the inversion of Bousquet et al. (2000) which have similar patterns of uptake in El Niño years as seen in our TCM results, although the release in 1983 is greater in the TCMs, and the uptake in 1989 is also larger in the TCMs. The magnitude of variability in the results of Bousquet et al. is $\pm 2 \text{ Gt C a}^{-1}$ compared with the $\pm 3 \text{ Gt C a}^{-1}$ for LPJ and IBIS, $\pm 2 \text{ Gt C a}^{-1}$ for HRBM and $\pm 1.5 \text{ Gt C a}^{-1}$ for TEM.

Figure 7

To investigate how the modeled source anomalies compare with the atmospheric CO₂ records we have calculated concentration anomalies for both the observations and simulated concentrations by again taking a 12 month running mean through the

detrended time series. An example of the concentration anomalies is shown in Figure 8 which shows the 12 month running mean for the observations and LPJ simulations at Mauna Loa in the left panel, and the modeled anomalies plotted against the observed anomalies with a linear regression in the right panel. In this example the observations show large negative anomalies for 1983 and 1987, which may appear to disagree with the source anomalies in Figure 7 which have positive anomalies in these years. In fact it should be expected that the concentration anomalies will lag the source anomalies as the concentration anomaly will continue to grow as long as the source anomaly is positive, such that the concentration anomaly will peak when the source anomaly returns to zero. On top of this, there will also be a lag between the source and concentrations due to the transport time.

Figure 8

Figure 9 shows the slope and correlation coefficients for linear regressions between the observed and modeled anomalies for each model for each station. The models all have slopes of less than 0.4. The low values are largely due to the failure of the TCM-MATCH simulations to model the large negative anomalies in the El Niño years which exist at many of the observing sites. The reason for not modeling these excursions may be because they result from a decrease in the tropical ocean source during the early phase on El Niño when the coastal upwelling is suppressed, and we have not considered the inter-annual variability in the ocean fluxes in this study. In the Mauna Loa example in Figure 8 the model underestimates the 1983 and 1987 anomalies by around 0.4 ppmv, which is equivalent to a global source of about 1 GtC, which is about the annual tropical release in our ocean simulation. If a significant portion of this release is suppressed in El

Figure 9

Niño years this could explain our low regression slope values. In general HRBM and TEM show lower slopes and correlations than IBIS and LPJ, indicating that IBIS and LPJ have more inter-annual variability than HRBM and TEM. IBIS and LPJ also tend to have higher correlations. On average the correlation required for significance at the 90% confidence limit (as determined by the random phase test of Ebisuzaki 1997) is 0.43, which is satisfied at most stations by IBIS and LPJ, but very few by TEM and HRBM. The models show lower correlations at KEY, NWR and CMO. It could be that oceans or transport play a more important role at these sites, or that the land areas of the northern mid-latitudes is the key region where the variability in the TCMs needs to be addressed.

5. Discussion

This study has evaluated the seasonal cycle and inter-annual variability from simulations of process-based TCMs. These analyses indicate that the models differ in their abilities to simulate these aspects of the global carbon cycle. No model consistently performs better than the others, suggesting that all four require further development. Below, we discuss the results of these analyses, point out some of the limitations of the study, and identify further research required to improved our understanding of the role of terrestrial processes in the global carbon cycle.

5.1. Average seasonal cycle

The comparison of the average atmospheric seasonal cycles for each of the TCM transport runs with the observations indicates that the models generally simulate the characteristics of the seasonal cycle, including the spatial trends in amplitude and phasing, although TEM and HRBM tend to perform better than LPJ and IBIS. These results, which were derived from the transient applications of the TCMs, are consistent with the results from other studies in which TCMs were run in equilibrium mode (see Heimann et al. 1998, Nemry et al. 1999, McGuire et al. 2000). The average seasonal cycle produced from the biosphere models for stations north of the tropics generally has too small a seasonal cycle, and the minimum concentration occurs too early in the year. Heimann et al. (1998) suggested that the biosphere models may overestimate the seasonal cycle of R_H . Reducing R_H during the growing season may allow NPP to dominate longer and thus correct the early phase and increase the seasonal amplitude. McGuire et al. (2000) demonstrated a possible mechanism to do this. By taking into account the thermal insulation effects of snow cover, this caused winter R_H to increase, while summer R_H decreased as the soil carbon pool was depleted. The overall effect was to increase the amplitude of the seasonal cycle of carbon uptake which led to an increase in the amplitude of the simulated atmospheric seasonal cycle of about 2 ppmv. As the TCMs tend to simulate the uptake of carbon too early in the spring, a better consideration of freeze-thaw dynamics by the models in high latitudes may improve the simulation of the average seasonal cycle at northern monitoring stations.

One region of disagreement between observations and the model simulations is in the tropics at ASC and SMO where there is substantial inter-annual variability in the observations and little agreement among the models. Being close to the equator, these sites are sensitive to both the Northern and Southern Hemisphere fluxes and the vigorous vertical transport in the tropics due to the large amount of convective activity. Also, fire, which has not been considered in the seasonal cycle here, may play an important part in the seasonal cycle in this region (Wittenberg et al. 1998, Iacobellis et al. 1994). The transport in the tropics has a large vertical component compared with other regions, as so much of the signal from the biosphere in this region is transported away from the surface and mixed with the upper troposphere. Thus, the impact of the region at the surface observing sites is relatively small. This is apparent when we consider that in the tropics the TCMs exhibit some of the strongest seasonality in the fluxes due to strong seasonality in rainfall (Tian et al., 1998, 2000), but the simulated atmospheric CO₂ concentrations in the tropics show relatively small amplitudes in the seasonal cycle. This suggests that the combination of atmospheric measurements with a transport model is not the ideal way to evaluate the TCMs in the tropics. An alternative method for evaluating the TCMs in this region may be aircraft measurements of CO₂, or column integrals from satellite measurements (Rayner et al. 2001)

5.2. Trends and inter-annual variability

The key advance in this study is the evaluation of the inter-annual variability from the TCMs. In general, the simulated trends in the amplitude of the seasonal cycle

are within the uncertainties of the observed trends. There is large uncertainty in the observed trends because of the short period we examine is, there is large uncertainty in the observed trends. The analysis of the TCM sources shows that CO₂ fertilization increased the amplitude, although the models vary in the magnitude, that climate variability acted to decrease the amplitude trend, and that land-use change in the northern mid-latitudes is probably the most important factor in driving the amplitude increase. This is an important result, as the effect of land-use change in the mid-latitudes is expected to decrease with time as the regrowth of the previously deforested areas is completed (Pacala et al. 2000).

The global source anomalies compare well with the most recent estimates from atmospheric inversions (Rayner et al. 2000, Bousquet et al. 2001), both in magnitude and phase. In fact, LPJ and IBIS tend to show more variability than the inversions. The TCMs show that most of the global variability comes from the tropics, and that the inter-annual variability is drive almost entirely by the climate variability. The comparison of inter-annual variability between the observed and simulated concentration anomalies suggests that the models produce considerably less variability than the observations, but that a significant factor in the underestimation is that the negative concentration anomalies associated with El Niño are not well modeled, and that the ocean inter-annual variability could explain much of the discrepancy. It is also possible that that the TCMs are not sensitive enough to inter-annual variability in the driving data (which is dominated by climate variability), or that the inter-annual variability in the transport is an important factor.

5.3. Limitations and next steps

The flask sampling network we have used employs a selection criteria to reduce the influence of local sources which are difficult to model with our relatively coarse resolution. Also, the frequency of measurements is much less than the sampling in the transport model. These factors may lead to biases, the consideration of which are beyond the scope of this paper. It is important to keep in mind that this evaluation is also subject to uncertainty as the atmospheric data set is also influenced by small scale processes which are not represented in our transport model.

The transport model also represents a source of uncertainty. The TransCom experiments have shown a range of transport behavior from different models. MATCH has been shown to be comparable with other widely used models, and its relatively fine resolution compared with GISS and TM2 models and sophisticated vertical parameterizations should allow for more accurate modeling of the transport. Evaluation and improvement of transport models is required for making progress in evaluating and improving TCMs. A potentially significant shortcoming of this study is that we have used model derived winds for our simulations. Clearly a useful experiment would be to repeat the transport runs using analysed winds fields, such as the National Center for Environmental Prediction Global Reanalyses (Kalnay et al. 1996).

The observing network used in this study is mostly in the Western Hemisphere, and so the fluxes from the Eastern Hemisphere biosphere are not well represented in the atmospheric data. There are currently a few stations in Asia, with GLOBALVIEW

sites in Korea, Mongolia and the South China Sea, but the sites have records starting in the early 1990s and are not appropriate for studying the 1980s. Once we are able to extend the simulations of the TCMs through the 1990s, the number of CO₂ observations available will increase substantially and we will be able to evaluate a much larger spatial domain. The longer time period will also allow a more rigorous investigation of the trends and inter-annual variability in the observed atmospheric seasonal cycle and the seasonal cycle simulated by the TCMs.

6. Conclusions

The key findings in this study are that when fluxes from four TCMs are run through an atmospheric transport model, they are able to reproduce the main features of the atmospheric seasonal cycle. Overall, the seasonal cycle from the TCMs tends to underestimate the amplitude and has the phase around 1 month early in the northern hemisphere. The main advance of this study has been to examine the contributions of the modeled terrestrial inter-annual variability. The key process driving the increase in the seasonal amplitude of atmospheric CO₂ is mid-latitude land-use change. The modeled concentration anomalies are shown to be driven by climate variability, but reproduce less than half of the observed variability, suggesting that either the ocean variability is a key factor, that models are not sensitive enough to the driving data, or that inter-annual variability in transport is also an important factor.

References

- Andres, R. J., G. Marland, I. Fung and E. Matthews, A $1^\circ \times 1^\circ$ distribution of carbon dioxide emissions from fossil fuel consumption and cement manufacture, 1950-1990, *Glob. Biogeochem. Cyc.*, *10*, 419-429, 1996.
- Battle, M., M. L. Bender, P. P. Tans, J. W. C. White, J. T. Ellis, T. Conway and R. J. Francey, Global carbon sinks and their variability inferred from atmospheric O_2 and $\delta^{13}C$, *Science*, *287*, 2467-2470, 2000.
- Bolin, B., R. Sukumar, P. Ciais, W. P. Cramer, P. Jarvis, H. Kheshgi, C. Nobre, S. Semenov and W. Steffen, Global Perspective. *IPCC Special Report on Land Use, Land-Use Change And Forestry*, R. T. Watson, I. R. Noble, B. Bolin, N. H. Ravinrdranath, D. J. Verardo and D. J. Dokken (Eds), Cambridge University Press, pp 23-51, 2000.
- Bousquet, P., P. Ciais, P. Peylin, M. Ramonet and P. Monfray, Inverse modeling of annual atmospheric CO_2 sources and sinks. 1. Method and control inversion, *J. Geophys. Res.*, *104*, 26161-26178, 1999.
- Bousquet, P., P. Peylin, C. LeQuéré, P. Friedlingstein and P. P. Tans, Regional changes in carbon dioxide fluxes over land and oceans since 1980, *Science*, *290*, 1342-1346, 2000.
- Chapin, F. S., S. A. Zimov, G. R. Shaver and S. E. Hobbie, CO_2 fluctuations at high latitudes, *Nature*, *383*, 585-586, 1996.
- Ciais, P., P. P. Tans, J. W. C. White, M. Trolier, R. J. Francey, J. A. Berry, D. R.

- Randall, P. J. Sellers, J. G. Collatz and D. S. Schimel, Partitioning of ocean and land uptake of CO₂ as inferred by $\delta^{13}\text{C}$ measurements from the NOAA Climate Monitoring and Diagnostics Laboratory Global Air Sampling Network, *J. Geophys. Res.*, *100*, 5051-5070, 1995.
- Cramer, W., D. Kicklighter, A. Bondeau, B. Moore, G. Churkina, B. Nemry, A. Ruimy, A. L. Schloss and N. Participants, Comparing global models of terrestrial net primary productivity (NPP): Overview and key results, *Global Change Biology*, *5*, 1-15, 1999.
- Dargaville, R. J., S. Doney, L. Kergoat, I. Y. Fung and D. S. Schimel, Covariance between inter-annual variability in CO₂ sources and atmospheric transport. *Abstracts volume, American Geophysical Union Fall Meeting, 2000, San Francisco, California*. AGU, 2000, pp 275.
- Denning, A. S., I. Y. Fung and D. A. Randall, Latitudinal gradient of atmospheric CO₂ due to seasonal exchange with land biota, *Nature*, *376*, 240-243, 1995.
- Doney, S. C., K. Lindsay and J. K. Moore, Global ocean carbon cycle modeling. *JGOFs/IGBP Synthesis*, M. Fasham (Ed), NCAR?, pp ???, 2001. In prep.
- Ebisuzaki, W., A method to estimate the statistical significance of a correlation when the data are serially correlated, *J. Climate*, *10*, 2147-2153, 1997.
- Enting, I. G., On the use of smoothing splines to filter CO₂ data, *J. Geophys. Res.*, *92D*, 10977-10984, 1987.
- Erickson III, D. J., P. J. Rasch, P. P. Tans, P. Friedlingstein, P. Ciais, E. Maier-Reimer,

- K. Six, C. A. Fischer and S. Walters, The seasonal cycle of atmospheric CO₂: A study based on the NCAR Community Climate Model (CCM2), *J. Geophys. Res.*, *101*, 15079-15097, 1996.
- Esser, G., J. Hoffstadt, F. Mack and U. Wittenberg, High resolution biosphere model (HRBM) - Documentation model version 3.00.00. *Mitteilungen aus dem Institut für Pflanzenökologie der Justus-Liebig-Universität Giessen*, G. Esser (Ed), Vol. 2, Giessen, Germany, pp 70, 1994.
- Fan, S., M. Gloor, J. Mahlman, S. Pacala, J. Sarmiento, T. Takahashi and P. Tans, A Large Terrestrial Carbon Sink in North America Implied by Atmospheric and Oceanic CO₂ Data and Models, *Science*, *282*, 442-446, 1998.
- Feely, R. A., R. Wanninkhof, T. Takahashi and P. Tans, Influence of El Nino on the equatorial Pacific contribution to atmospheric CO₂ accumulation, *Nature*, *398*, 597-601, 1999.
- Foley, J. A., I. C. Prentice, N. Ramankutty, S. Levis, D. Pollard, S. Sitch and A. Haxeltine, An integrated biosphere model of land surface processes, terrestrial carbon balance, and vegetation dynamics, *Global Biogeochem. Cycles*, *10*, 603-628, 1996.
- GLOBALVIEW-CO₂. Cooperative Atmospheric Data Integration Project - Carbon Dioxide. CD-ROM, NOAA/CMDL, Boulder, Colorado, 1999. [Also available on Internet via anonymous FTP to ftp.cmdl.noaa.gov, Path: ccg/co2/GLOBALVIEW].

- Hack, J. J., Parameterization of moist convection in the NCAR Community Climate Model, CCM2, *J. Geophys. Res.*, *99*, 5551-5568, 1994.
- Heimann, M., G. Esser, A. Haxeltine, J. Kaduk, D. Kicklighter, W. Knorr, G. Kohlmaier, A. McGuire, J. Melillo, B. Moore, R. Otto, I. Prentice, W. Sauf, A. Schloss, S. Sitch, U. Wittenberg and G. Wurth, Evaluation of terrestrial carbon cycle models through simulations of the seasonal cycle of atmospheric CO₂: First results of a model intercomparison study, *Glob. Biogeochem. Cyc.*, *12*, 1-24, 1998.
- Holtslag, A. A. M., and B. A. Boville, Local Versus Nonlocal Boundary-Layer Diffusion in a Global Climate Model, *J. Clim.*, *6*, 1825-1842, 1993.
- Houghton, R. A., The annual net flux of carbon to the atmosphere from changes in land use 1850-1990, *Tellus*, *51B*, 298-313, 1999.
- Hulme, M. A., 1951-80 global land precipitation climatology for the evaluation of General Circulation Models, *Climate Dynamics*, *7*, 57-72, 1992.
- Iacobellis, S. F., R. Frouin, H. Razafimpanilo, R. C. J. Somerville and S. C. Piper, North African savanna fires and atmospheric carbon dioxide, *J. Geophys. Res.*, *99*, 8321-8334, 1994.
- Jones, P. D., Hemispheric surface air temperature variations; a reanalysis and an update to 1993, *J. Climate*, *7*, 1794-1802, 1994.
- Joos, F., R. Meyer, M. Bruno and M. Leuenberger, The variability in the carbon sinks as reconstructed for the last 1000 years, *Geophys. Res. Lett.*, *26*, 1437-1440, 1999.

- Kalnay, E., The NMC/NCAR 40-year reanalysis project, *Bull. Amer. Meteor. Soc.*, *77*, 437-471, 1996.
- Keeling, C. D., T. P. Whorf, M. Wahlen and J. van der Plicht, Interannual extremes in the rate of rise of atmospheric carbon dioxide since 1980, *Nature*, *375*, 666-670, 1995.
- Keeling, C. D., J. F. S. Chin and T. P. Whorf, Increased activity of northern vegetation inferred from atmospheric CO₂ measurements, *Nature*, *382*, 146-149, 1996.
- Kohlmaier, G. H., E. O. Sire, A. Janecek, C. D. Keeling, S. C. Piper and R. Revelle, Modeling the seasonal contribution of a CO₂ fertilization effect of the terrestrial vegetation to the amplitude increase in atmospheric CO₂ at Mauna Loa observatory, *Tellus*, *41B*, 487-510, 1989.
- Lee, K., R. Wanninkhof, T. Takahashi, S. C. Doney and R. A. Feely, Low interannual variability in recent oceanic uptake of atmospheric carbon dioxide, *Nature*, *396*, 155-159, 1997.
- LeQuéré, C., J. C. Orr, P. Monfray, O. Aumont and G. Madec, Interannual variability of the oceanic sink of CO₂ from 1979 through 1997, *Global. Biogeochem. Cycles*, *14*, 1247-1266, 2000.
- Mahowald, N. M., P. J. Rasch, B. E. Eaton, S. Whittlestone and R. G. Prinn, Transport of ²²²radon to the remote troposphere using the Model of Atmospheric Transport and Chemistry and assimilated winds from ECMWF and the National Center for Environmental Prediction/NCAR, *J. Geophys. Res.*, *102*, 28139-28151, 1997.

Marland, G., T. Boden and R. J. Andres, Global, Regional, and National CO₂-Emissions.

Trends: A Compendium of Data on Global Change, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. DoE, Oak Ridge, Tenn, U.S.A., 2000.

Masarie, K. A., and P. P. Tans, Extension and integration of atmospheric carbon dioxide data into a globally consistent measurement record, *J. Geophys. Res.*, *100*, 11593-11610, 1995.

McGuire, A. D., J. M. Melillo, J. T. Randerson, W. J. Parton, M. Heimann, R. A. Meier, J. S. Clein, D. W. Kicklighter and W. Sauf, Modeling the effect of snowpack on the heterotrophic respiration across northern temperate and high latitude regions: Comparison with measurements of atmospheric carbon dioxide in high latitudes, *Biogeochemistry*, *48*, 91-114, 2000.

McGuire, A., S. Sitch, R. Dargaville, G. Esser, J. Foley, M. Heimann, F. Joos, J. Kaplan, D. W. Kicklighter, R. A. Meier, J. M. Mellilo, B. Moore, I. C. Prentice, N. Ramankutty, T. Reichenau, A. Schloss, H. Tian and U. Wittenberg, Carbon balance of the terrestrial biosphere in the twentieth century: Analysis of CO₂, climate and land-use effects with four processed-based models, *Global Biogeochem. Cycles*, *15*, 183-203, 2001.

Nemry, B., L. Francois, J. C. Gerard, A. Bondeau, M. Heimann and PNMIG, Comparing global models of terrestrial net primary productivity (NPP): analysis of the seasonal atmospheric CO₂ signal, *Global Change Biology*, *5*, 65-76, 1999.

Pacala, S. W., G. C. Hurtt, R. A. Houghton, R. A. Birdsey, E. T. Sundquist, D. Baker, P. Peylin, P. Moorcroft, J. Caspersen, M. E. Harmon, S. M. Fan, J. L. Sarmiento, C. Goodale and C. B. Field, Carbon storage in the US caused by land use change: convergence of estimates from inventories of ecosystems and inversions of atmospheric data. *Abstracts volume, American Geophysical Union Fall Meeting, 2000, San Francisco, California*. AGU, 2000, pp 275.

Ramankutty, N., and J. A. Foley, Characterizing patterns of global landuse: An analysis of global croplands data, *Global Biogeochem. Cycles*, *12*, 667-685, 1998.

Randerson, J. T., M. V. Thompson, T. J. Conway, I. Y. Fung and C. B. Field, The contribution of terrestrial sources and sink to trends in the seasonal cycle of atmospheric carbon dioxide, *Global. Biogeochem. Cycles*, *11*, 535-560, 1997.

Randerson, J. T., C. B. Field, I. Y. Fung and P. P. Tans, Increases in early season ecosystem uptake explain recent changes in the seasonal cycle of atmospheric CO₂ at high northern latitudes, *Geophys. Res. Lett.*, *26*, 2765-2768, 1999.

Rasch, P., N. M. Mahowald and B. E. Eaton, Representations of transport, convection and the hydrologic cycle in chemical transport models: Implications for the modeling of short lived and soluble species, *J. Geophys. Res.*, *102*, 28127-28138, 1997.

Rayner, P. J., and D. M. O'Brien, The utility of remotely sensed CO₂ concentration data in surface inversions, *Geophys. Res. Lett.*, *28*, 175-178, 2001.

Rayner, P. J., I. G. Enting, R. J. Francey and R. L. Langenfelds, Reconstructing the

- recent carbon cycle from atmospheric CO₂, δ¹³C and O₂/N₂ observations, *Tellus*, 51B, 213-232, 1999.
- Schimel, D., J. Melillo, H. Tian, A. D. McGuire, D. Kicklighter, T. Kittel, N. Rosenbloom, S. Running, P. Thornton, D. Ojima, W. Parton, R. Kelly, M. Sykes, R. Neilson, B. Rizzo and L. Pitelka, Carbon storage by natural and agricultural ecosystems of the US 1980-1993, *Science*, 287, 2004-2006, 2000.
- Sitch, S., I. C. Prentice, B. Smith and J. O. Kaplan, A coupled model of vegetation dynamics and the terrestrial carbon cycle, in prep.
- Tian, H., J. M. Melillo, D. W. Kicklighter, A. D. McGuire, J. V. K. Helfrich, B. Moore and C. Vörösmarty, Effect of interannual climate variability on carbon storage in Amazonian ecosystems, *Nature*, 396, 664-667, 1998.
- Tian, H., J. M. Melillo, D. W. Kicklighter and A. D. McGuire, The sensitivity of terrestrial carbon storage to historical climate variability and atmospheric CO₂ on the United States, *Tellus*, 51B, 414-452, 1999.
- Tian, H., J. M. Melillo, D. W. Kicklighter, A. D. McGuire, J. Helfrich, B. Moore and C. J. Vörösmarty, Climatic and biotic controls on annual carbon storage in Amazonian ecosystems, *Global Ecology and Biogeography*, 9, 315-336, 2000.
- VEMAP, Vegetation/ecosystem modeling and analysis project: Comparing biogeography and biogeochemistry models in a continental-scale study of terrestrial ecosystem responses to climate change and CO₂ double, *Global Biogeochem. Cycles*, 9, 407-437, 1995.

Waugh, D. W., Three-dimensional simulations of long-lived tracers using winds from

MACCM2, *J. Geophys. Res.*, *102*, 21493-21513, 1997.

Wittenberg, U., M. Heimann, G. Esser, A. D. McGuire and W. Sauf, On the influence of

biomass burning on the seasonal CO₂ signal as observed at monitoring stations,

Glob. Biogeochem. Cycles, *12*, 531-544, 1998.

Zimov, S. A., S. P. Davidov, G. M. Zimova, A. I. Davidova, F. S. Chapin, M. C. Chapin

and J. F. Reynolds, Contribution of disturbance to increasing seasonal amplitude

of atmospheric CO₂, *Science*, *284*, 1973-1976, 1999.

J. S. Clein, R. Dargaville, R. A. Meier, Institute of Arctic Biology, PO Box 700,
University of Alaska, Fairbanks 99775 AK, (email fnjsc4@uaf.edu, rjd@ucar.edu,
fnram2@uaf.edu)

G. Esser, T. Reichenau, U. Wittenberg, Institut für Pflanzenökologie, Justus-Liebig-
Universität, Giessen, Germany (email esser@bio.uni-giessen.de, tim.reichenau@bio.uni-
giessen.de, uwe.wittenberg@merck.de)

J. Foley, N. Ramankutty, Climate, People and Environment Program, Institute
for Environmental Studies, University of Wisconsin, Madison, Wisconsin (email
jfoley@facstaff.wisc.edu, nramanku@students.wisc.edu)

M. Heimann, J. Kaplan, I. C. Prentice, Max-Planck Institut für Biogeochemie, Jena,
Germany (email martin.heimann@bgc-jena.mpg.de, colin.prentice@bgc-jena.mpg.de,
jkaplan@bgc-jena.mpg.de)

F. Joos, Physics Institute, University of Bern, Bern, Switzerland (email joos@climate.unibe.ch)

D. W. Kicklighter, J. M. Melillo, H. Tian, The Ecosystems Marine Biological Laboratory, Woods Hole, Massachusetts (email dkick@mbl.edu, jmelillo@mbl.edu htian@mbl.edu)

A. D. McGuire U. S. Geological Survey, Alaska Cooperative Fish and Wildlife Research Unit, University of Alaska, Fairbanks 99775 AK (email ffadm@uaf.edu)

B. Moore III, A. Schloss, Complex Systems Research Center, Institute for the Study of Earth, Oceans and Space, University of New Hampshire. Durham, New Hampshire (email b.moore@unh.edu, annette.schloss@unh.edu)

S. Sitch, Potsdam Institut for Climate Impact Research, Potsdam, Germany (email sitch@pik-potsdam.de)

L. J. Williams, Electric Power Research Institute, Palo Alto, California (email ljwillia@epri.com)

Received ???; revised ???; accepted ???.

¹Institute of Arctic Biology, University of Alaska, Fairbanks, Alaska

²Max-Plank-Institut für Biogeochemie, Jena, Germany

³US Geological Survey, Alaska Cooperative Fish and Wildlife Research Unit, University of Alaska, Fairbanks, Alaska

⁴The Ecosystem Center, Marine Biological Laboratory, Woods Hole, Massachusetts

⁵Physics Institute, University of Bern, Bern, Switzerland

⁶Institut für Pflanzenökologie, Justus-Liebig-Universität, Giessen, Germany

⁷Climate, People and Environment Program, Institute for Environmental Studies,
University of Wisconsin, Madison, Wisconsin

⁸Complex Systems Research Center, Institute for the Study of Earth, Oceans and
Space, University of New Hampshire. Durham, New Hampshire

⁹Potsdam Institute for Climate Impact Research, Potsdam, Germany

¹⁰Electric Power Research Institute, Palo Alto, California

¹¹Now at National Center for Atmospheric Research, Boulder, Colorado

Figure 1. Location of 15 NOAA/CMDL CO₂ monitoring sites. Note that Mauna Loa and Kumukahi share almost the same latitude and longitude, but that Kumukahi is at sea level while Mauna Loa is at 3400m. See Figure 3 for abbreviation expansions.

Figure 2. Average seasonal cycle aggregated over 30° latitude bands for the period 1980 to 1991 for the four biosphere models HRBM, IBIS, LPJ and TEM. The southern extratropical regions are not shown due to the small amount of biosphere in these regions. Note that the first six months of the annual cycle have been repeated.

Figure 3. Average detrended seasonal cycle at 15 NOAA/CMDL monitoring sites for the GLOBALVIEW (1999) observations, and transport model runs with sources from HRBM, IBIS, LPJ and TEM. The pattern for each TCM result includes the contribution from the modeled fossil and ocean sources, which are also plotted. The individual captions detail the names and locations (latitude and longitude) of the stations and the three-letter abbreviations used in the text.

Figure 4. Seasonal cycle analysis at 15 NOAA/CMDL stations for the transport run using the S3 fluxes from the four TCMs. (a) Amplitude of the seasonal cycle (b) timing of minimum concentrations and (c) timing of the maximum concentrations. Note that the stations between 0 and 30°S have been equally spaced with latitude to make it easier to distinguish the different values. Each of the models is shown by a point symbol, while the observations are shown as a shaded area showing the range of one standard deviation about the mean.

Figure 5. Changes in amplitude of the seasonal cycle of TCM sources aggregated over 30° latitude bands for each of the three scenarios: 1: increasing CO₂, 2: increasing CO₂ and climate variability, and 3: increasing CO₂, climate variability and land-use change.

Figure 6. The trend in the seasonal cycle amplitude at the 15 NOAA/CMDL sites for each of the three TCM scenarios for (a) HRBM, (b) IBIS, (c) LPJ and (d) TEM. Trends were calculated by a linear regression through the time series of seasonal amplitudes. Shaded areas represent the observed gradient range as determined by the 90% confidence limits of the regression.

Figure 7. Source anomalies for each of the TCMs aggregated over 30° latitude bands, and the global sum. Anomalies were calculated by taking the 12 month running mean through the monthly source values. The bold solid line in the global panel is from the Bousquet et al. (2000) inversion.

Figure 8. Example of the time series of the concentration anomalies and the regression. This example shows the comparison between the LPJ model and observations for Mauna Loa.

Figure 9. Results from regression analysis of simulated and observed concentration anomalies derived by taking the running mean through the trended timeseries. (a) is the slope of the regression and (b) is the correlation of the simulated against observed anomalies. The dashed line shows the correlation requirement for 90% significance.

Acknowledgments. This work was conducted as part of the Carbon Cycle Model Linkage Project with support from the Electric Power Research Institute, from the Land Cover Land Use Change Program of the National Aeronautics and Space Administration, and from the National Science Foundation. This work was supported in part by a grant of HPC time from the Arctic Region Supercomputing Center. The authors would like to thank Philippe Bousquet et al. for their data from the 2000 Science paper and Scott Doney for helpful comments on the manuscript.

















